



**QUEEN'S
UNIVERSITY
BELFAST**

Seasonality of pCO₂ in a hard-water lake of the northern Great Plains: The legacy effects of climate and limnological conditions over 36 years

Finlay, K., Vogt, R. J., Simpson, G. L., & Leavitt, P. R. (2019). Seasonality of pCO₂ in a hard-water lake of the northern Great Plains: The legacy effects of climate and limnological conditions over 36 years. *Limnology and Oceanography*, 64, S118-S129. <https://doi.org/10.1002/lno.11113>

Published in:
Limnology and Oceanography

Document Version:
Publisher's PDF, also known as Version of record

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

Copyright 2019 the authors.
This is an open access article published under a Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution and reproduction in any medium, provided the author and source are cited.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Seasonality of pCO₂ in a hard-water lake of the northern Great Plains: The legacy effects of climate and limnological conditions over 36 years

Kerri Finlay ^{1,*} Richard J. Vogt,² Gavin L. Simpson ^{1,3} Peter R. Leavitt ^{1,3,4}

¹Department of Biology, University of Regina, Regina, Saskatchewan, Canada

²Department of Biology, York University, Toronto, Ontario, Canada

³Institute of Environmental Change and Society, University of Regina, Regina, Saskatchewan, Canada

⁴Institute for Global Food Security, Queen's University Belfast, Belfast, County Antrim, United Kingdom

Abstract

Biogeochemical processes are active year-round in ice-covered lakes, such that processes in one season can affect limnological conditions in subsequent seasons. However, the extent and nature of these legacy effects are poorly understood, particularly for the CO₂ content of lakes and when considering gas exchange with the atmosphere. Here, we used a unique 36-yr dataset of weekly limnological measurements of Buffalo Pound Lake in the northern Great Plains to assess seasonal changes in CO₂ concentration and flux and determine how dependent lake pCO₂ is on limnological conditions of previous seasons. We found that the lake was a net source of CO₂ to the atmosphere (mean 18.5 ± 7.4 mol CO₂ m⁻² yr⁻¹), with spring potentially accounting for the majority (~ 64% ± 20%) of CO₂ efflux, assuming ice in spring was permeable to gas exchange (32.9% ± 19.8% if not). Analysis with generalized additive models (GAMs) demonstrated that current and antecedent seasonal conditions combined to explain 72.6% of deviance in spring pCO₂, but that the strength of model predictions and the importance of antecedent conditions diminished in GAMs of summer (43.6%) and fall (23.3%) CO₂ levels. This research suggests that pCO₂ is regulated by a combination of coeval and historical environmental conditions and shows that quantification of seasonal and annual fluxes requires a mechanistic understanding of the legacy effects of preceding time intervals.

It is now well established that inland waters contribute significantly to the global carbon budget (Cole et al. 2007; Prairie 2008; Tranvik et al. 2009), although many questions remain about the factors regulating variability in water-column pCO₂ at broad spatial and temporal scales. One such uncertainty relates to the legacy effects of antecedent water-column conditions on current ecosystem function. For example, biogeochemical cycling under ice can substantially alter the abundance and chemical form of macronutrients in spring (Kratz et al. 1987; Hampton et al. 2017) and, in the case of carbon (C),

substantially increase CO₂ concentrations under ice (Kratz et al. 1987; Finlay et al. 2015). Additionally, although spring CO₂ flux has been shown to contribute significantly to total annual CO₂ flux in many lakes (Maberly 1996; Striegl et al. 2001; Ducharme-Riel et al. 2015), relatively few measurements of pCO₂ are available for shoulder seasons of summer, owing to logistical issues related to sampling during ice melt and formation. Given that lake pCO₂ is frequently elevated in spring and fall seasons relative to summer (Baehr and DeGrandpre 2002; Denfeld et al. 2015), it is important to better understand the magnitude and drivers of seasonal contributions to annual CO₂ fluxes to improve estimates of the role lakes in the global C cycle.

Seasonal variation in water-column pCO₂ in boreal lakes frequently follows predictable annual patterns of change in metabolic processes, particularly in ice-covered dimictic systems. In these lakes, CO₂ accumulates under ice in winter (Baehr and DeGrandpre 2002; Denfeld et al. 2015), causing a large efflux of CO₂ in spring when the ice melts and the water column circulates (overturn). pCO₂ levels are reduced in summer when the water column is stable and primary production

*Correspondence: kerri.finlay@uregina.ca

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Additional Supporting Information may be found in the online version of this article.

Special Issue: Long-term Perspectives in Aquatic Research.

Edited by: Stephanie Hampton, Matthew Church, John Melack and Mark Scheuerell.

increases, while $p\text{CO}_2$ often increases during fall when CO_2 from respired organic matter in the hypolimnion is mixed into the water column at fall overturn (Maberly 1996; Dillon and Molot 1997; Anderson et al. 1999; Baehr and DeGrandpre 2004; Ducharme-Riel et al. 2015). Deviations from this pattern can occur due to local variation in meteorological conditions (wind, atmospheric pressure, and storm runoff), which affect lake stratification and gas solubility (Vachon and del Giorgio 2014) or which introduce labile allochthonous organic matter into the lake (López-Bellido et al. 2012).

Less is known about seasonal patterns of $p\text{CO}_2$ in hard water and saline lakes that account for nearly half of continental surface waters (Hammer 1986). In these hard-water systems, variation in pH, groundwater inputs, and calcite precipitation can uncouple lake $p\text{CO}_2$ from metabolically regulated processes (Striegl and Michmerhuizen 1998; Finlay et al. 2009; Stets et al. 2009). Moreover, the magnitude of atmospheric CO_2 exchange in spring and fall has not been widely quantified in the shallow polymictic lakes common in agricultural lowlands, but where summer CO_2 effluxes can be much less than that seen in dimictic lakes (Finlay et al. 2009, 2015). In particular, hypolimnetic CO_2 accumulation should be relatively low in the absence of persistent thermal stratification, whereas frequent lake mixing should keep the vertical profiles of $p\text{CO}_2$ more uniform during the ice-free period (Anderson et al. 1999; Stets et al. 2009). Given the potential importance of such hard-water lakes in regulating atmospheric CO_2 exchange (Finlay et al. 2015) and the predominance of spring and fall CO_2 emissions in other lakes (Ducharme-Riel et al. 2015), further research is needed on the controls of seasonal and annual CO_2 content in polymictic hard-water lakes.

Under-ice processes can be influential for many biogeochemical cycles, including that of carbon (Kratz et al. 1987), and can thus result in legacy effects where antecedent conditions propagate into subsequent seasons (Meding and Jackson 2003; Hampton et al. 2017; Powers et al. 2017). For example, respiration rates can be high under ice (Denfeld et al. 2015), particularly near the sediments where the temperature tends to be warmer than in surface waters and where organic matter accumulation is high (Wetzel 2001). Photosynthesis can also be an important control of CO_2 immediately under ice when snow cover is limited (Baehr and DeGrandpre 2002, 2004; Pernica et al. 2017). As a result, the quantity of CO_2 accumulated under ice can be a function of the duration of ice cover (Finlay et al. 2015), the availability of nutrients and light for photosynthesis (Baehr and DeGrandpre 2002, 2004; Salmi and Salonen 2016; Pernica et al. 2017), and the quantity and quality of organic matter available for mineralization (Wetzel 2001; Hampton et al. 2017). Similarly, respiratory consumption of O_2 under ice is dependent on winter conditions, as well as previous seasons' primary production (Meding and Jackson 2003). However, while biogeochemical processes during winter conditions can affect limnological conditions in spring, it is less clear how these processes affect CO_2 flux at

ice-off, or whether winter legacy effects continue through summer and fall.

Together, this evidence suggests that water-column $p\text{CO}_2$ at a given point in time is dependent on both present limnological conditions as well as those in preceding seasons. To evaluate this hypothesis, we used generalized additive models (GAMs) to quantify the magnitude and correlates of seasonal and annual CO_2 dynamics in a polymictic eutrophic hard-water lake that has been monitored year-round at weekly intervals for 36 yr. Our objectives were three-fold: (1) describe seasonal variation (spring, summer, and fall) in $p\text{CO}_2$ and potential atmospheric exchange in a polymictic lake; (2) quantify long-term (36 yr) trends in CO_2 dynamics and seasonality; and (3) evaluate the influence of antecedent environmental conditions (productivity and climate) on seasonal estimates of water-column $p\text{CO}_2$. We predicted that spring $p\text{CO}_2$ would be influenced strongly by factors controlling the supply of labile organic matter and the duration of ice cover (Meding and Jackson 2003; Finlay et al. 2015) but that atmospheric CO_2 exchange in summer and fall would reflect the increasing influence of coeval meteorological and limnological conditions (Gerten and Adrian 2000; Winder and Schindler 2004). By integrating seasonal change with the importance of legacy effects, we hope to improve predictions of how future climate change may affect the contribution of lakes to the global carbon budget.

Methods

Study site

Buffalo Pound Lake is a natural lake that was impounded in 1939 and 1952 by the damming of the outflow into the Qu'Appelle River in southern Saskatchewan, Canada (Hall et al. 1999). The lake is long and narrow (1 km by 29 km), with an average depth of 3 m. The shallow depth of the lake, combined with long fetch along the prevailing storm track, results in a polymictic system that only rarely establishes weak thermal stratification (Dröscher et al. 2009).

Buffalo Pound provides the drinking water supply for the cities of Moose Jaw (population 45,000) and Regina (population 216,000), has water levels managed by the Saskatchewan Water Security Agency, and is maintained in part by hydrologic transfer from the upstream Lake Diefenbaker reservoir (Hall et al. 1999). The lake receives runoff from a 3310 km² agricultural catchment area in which nutrient-rich soils favor high nutrient influx and eutrophic conditions. Although other high pH lakes in this region typically ingest CO_2 (Finlay et al. 2015), Buffalo Pound is typically oversaturated with CO_2 and exhibits net outgassing of CO_2 to the atmosphere over the last 20 yr (Finlay et al. 2015).

As the main urban drinking water supply, Buffalo Pound has been monitored on a weekly basis since 1979 for 65 water quality parameters. Raw water taken from an inflow pipe 1 m above the bottom of the lake, at 3 m depth, is pumped into the water treatment plant for analyses, treatment for human

use, and distribution. Measured parameters include physical (temperature), chemical (pH, nutrients, and major ions), and biological (chlorophyll *a* [Chl *a*], algae, and bacteria) properties. In this study, we used measured conductivity ($\mu\text{S cm}^{-1}$), bicarbonate and carbonate (mg L^{-1}), pH, and temperature ($^{\circ}\text{C}$) to calculate $p\text{CO}_2$ and potential CO_2 flux. Furthermore, we explored proxies of planktonic metabolism (Chl *a* and dissolved organic carbon [DOC]) and physicochemical processes (water temperature and ice-cover duration) as predictors of variation in $p\text{CO}_2$ and CO_2 fluxes across the 36-yr time series.

$p\text{CO}_2$ and CO_2 flux calculations

CO_2 concentration (μM), $p\text{CO}_2$ (μatm), and CO_2 flux ($\text{mmol CO}_2 \text{ m}^{-2} \text{ d}^{-1}$) were estimated for each sampling date from conductivity, water temperature, and pH measurements taken from the inflow water as described in Finlay et al. (2015). Dissolved inorganic carbon (DIC) concentrations were estimated using a previously derived relationship between measured DIC and conductivity for Buffalo Pound Lake ($r^2 = 0.98$, $p < 0.001$; Finlay et al. 2009). Given the elevated pH of the system (average pH during the open water period = 8.3), chemically enhanced C flux was calculated on each sampling date.

CO_2 flux was calculated as

$$\text{net daily CO}_2\text{flux} = \alpha k([\text{CO}_2]_{\text{lake}} - [\text{CO}_2]_{\text{sat}}),$$

where $[\text{CO}_2]_{\text{lake}}$ is the concentration of CO_2 in the water, $[\text{CO}_2]_{\text{sat}}$ is the concentration of CO_2 at equilibrium with the atmosphere, α is the chemical enhancement of CO_2 flux at high pH (Hoover and Berkshire 1969), and k is piston velocity (cm h^{-1}) as determined from Model B in Vachon and Prairie (2013), relating k to wind speed and lake surface area. Hourly wind speed as measured each day at 10-m height was collected from publicly available Environment Canada records for the city of Moose Jaw (Sta. 2967; <http://climate.weather.gc.ca/>). Flux was interpolated between time points by multiplying daily flux rates by 7 d to get a total potential flux for each of 52 weeks. The concentration of CO_2 at saturation with the atmosphere was taken as the global mean annual CO_2 concentration measured at Mauna Loa observatory.

Raw water was collected between 07:00 and 07:30 h, once each week, and thus diel variations in pH were not considered in this analysis. Although pH can vary considerably during the day (Maberly 1996), an evaluation of continuous sonde data (15 min resolution) from Buffalo Pound during summer 2014 suggested no systematic bias in $p\text{CO}_2$ estimates due to the use of morning pH measurements. Specifically, $p\text{CO}_2$ at 07:00 h was correlated positively with mean daily $p\text{CO}_2$ ($r = 0.44$, $p < 0.001$) and was not consistently elevated or depressed relative to the daily values. Similarly, the $p\text{CO}_2$ values calculated from water treatment plant samples should be elevated relative to those of surface waters (Finlay et al. 2015), as water was extracted 1 m above the bottom (2 m from surface) and is more affected by sedimentary

respiration and less by photosynthesis in this turbid system (mean summer Secchi depth = 1.1 m). However, given that the lake is polymictic year-round, we assumed that these surface-deep differences did not greatly influence our analysis of temporal variability and causal relationships, at least during the open water period (Finlay et al. 2015).

Ice-on and ice-off dates were provided by the Buffalo Pound water treatment plant and were determined as the day of year when the lake was 90% covered with ice (ice-on) or 90% ice free (for ice-off).

Definition of seasons

As the goal of this study was to quantify the magnitude of variation in seasonal $p\text{CO}_2$, identify potential controls thereof, and evaluate the importance of antecedent seasonal conditions on observed CO_2 , we needed to establish functional definitions to delineate seasons, which accounted for interannual variation in winter severity and ice-cover duration. Complete ice melt can take weeks and may vary with spring meteorology (Finlay et al. 2015), so we defined seasons based on a combination of potential gas exchange with the atmosphere (winter and spring) and consistent static calendar dates (summer and fall) that define when CO_2 concentrations were stable, as recommended by Anderson et al. (1999).

Herein, winter was defined as the period when the lake was completely covered with ice, and atmospheric gas exchange was negligible, beginning with the date of ice formation in the fall and continuing until the date of maximum modeled CO_2 concentration in Buffalo Pound (see below). The start of spring was defined as the date when CO_2 concentrations begin to decline, assuming that this pattern arises from loss of CO_2 to the atmosphere (see Discussion section), even if this occurred before the recorded ice-off date. Spring continued until the minimum CO_2 concentration was recorded (within 100 d of CO_2 maxima). Spring was further divided into two phases for CO_2 flux analyses: “potential spring,” which was defined as the period of time between maximum CO_2 concentration and ice-off date, and “open-water spring,” the period from documented ice-off until the minimum CO_2 concentration. Hereafter, “spring” refers to the period that includes both “potential” and “open-water” spring periods, and annual flux rates include “potential” spring, unless otherwise indicated. Summer was then calculated as the date after the spring CO_2 minimum continuing until 31 August of that year, whereas fall was defined as the period from 01 September until ice-on, as we have done previously (Finlay et al. 2015).

Statistical analyses

The time series of in situ $p\text{CO}_2$ was modeled using a GAM (Hastie & Tibshirani 1990; Wood et al 2016; Wood 2017; Simpson 2018) comprising terms to account for both within- and between-year variation in the time series. We chose to model the data using a GAM, because this approach better




Fig. 1. Total annual CO₂ flux in Buffalo Pound from 1979 to 2014, broken down by contribution by each open-water season. Spring is divided into “potential spring,” the period between maximum pCO₂ and the ice-off date (black bars), and “open-water spring,” the period between ice-off and minimum pCO₂ (dark gray bars). Summer and fall are represented by medium and light gray bars, respectively. Flux is calculated using summed calculated CO₂ flux based on weekly data, extrapolated to 7 d. Winter flux under ice was considered to be zero.

accounts for nonlinearity of trends relative to other protocols (e.g., Mann Kendall test) and GAMs uniquely allow us to estimate the magnitudes of within- and between-year trends in the data, derive secondary estimates from the model (e.g., the magnitude of efflux at ice-out), and quantify uncertainties. For example, the commonly used (seasonal) Mann Kendall test does not estimate the magnitude(s) of trends, tests only for monotonic trends (which were not indicated in preliminary data screening), and does not allow derivation of secondary estimates as above. Similarly, estimation of trends using parametric linear or generalized linear models would require us to a priori state the functional form of the within- and between-year trends in time series or perform model selection from among a set of complex polynomial models. Using GAMs, we avoid this subjective element of model specification by allowing the functional form of the trends to be determined from the data, whereas the use of splines avoids well-known bias issues at the ends of series that plague polynomial models. Details of candidate model selection and estimates are included in the Supporting Information.

To determine the start and end dates of spring, the best-fitting GAM for the pCO₂ time series was used to estimate annual mean differences between the minimum and maximum pCO₂ between days 50 (19 February) and 160 (~ 09 June). In this procedure, an estimate for the expected difference for each year can be derived by predicting daily pCO₂ for each day in the specified interval, finding the peak pCO₂ during the period, and calculating the difference between the two pCO₂ extremes. Of the 36 yr of data collection, 4 yr did not have a pronounced pCO₂ peak and those years were not included in subsequent statistical analyses. Uncertainty in the estimated pCO₂ trend was evaluated using 10,000 simulations

of the trend from the posterior distribution of the fitted GAM (details in Supporting Information).

Variables known from the literature to affect pCO₂ content of prairie hard-water lakes were selected a priori to develop individual GAMs for each season to predict pCO₂ in Buffalo Pound (Meding and Jackson 2003; Finlay et al. 2009, 2010). Specifically, spring CO₂ flux was expected to be dependent on ice-cover duration (longer ice cover resulting in greater accumulation of respired CO₂), and the productivity of the previous summer (providing the material for respiration over winter) approximated as Chl *a* (Meding and Jackson 2003; Finlay et al. 2015). In contrast, we expected that summer and fall pCO₂ would be more heavily dependent on pCO₂ in the previous season and coeval limnological conditions (Finlay et al. 2009, 2010). To test these hypotheses, we developed GAMs to evaluate the effects of coeval mean water temperature, Chl *a*, DOC, and ice-cover duration in the models and examined legacy effects on seasonal lake pCO₂ by including mean values from preceding seasons. Given that not all variables were measured in all years, direct comparisons of Akaike information criteria (AIC) were not appropriate for determining the best-fitting model for each season. Instead, we selected models that maximized deviance explained, adjusted *R*² (*R*_{adj}²), and sample size (*n*), in addition to a qualitative exploration of the model fits. More details of model selection are included in the Supporting Information.

GAMs were estimated using the *mgcv* package (version 1.8-22; Wood 2017), and graphics were plotted with package *ggplot2* (Wickham 2009) for R (version 3.5.1; R Core Team 2018). R scripts and data to reproduce the analysis can be found at <https://doi.org/10.5281/zenodo.2431144>.




Fig. 2. Seasonally averaged $p\text{CO}_2$ by year of study in Buffalo Pound. No directional trends were observed for any season over the 36-yr period.

Results

Estimation of $p\text{CO}_2$ from water chemistry suggested that Buffalo Pound Lake should outgas CO_2 during the open-water season of most years (Fig. 1). Estimates of total annual CO_2 flux (including potential spring flux) ranged from a minimum of $4.36 \text{ mol m}^{-2} \text{ yr}^{-1}$ in 1988 to a maximum of $41.97 \text{ mol m}^{-2} \text{ yr}^{-1}$ in 1992, with a mean ($\pm \text{SD}$) annual flux rate of $18.53 \pm 7.38 \text{ mol m}^{-2} \text{ yr}^{-1}$. Instantaneous CO_2 fluxes ranged dramatically from an efflux of $886.8 \text{ mmol m}^{-2} \text{ d}^{-1}$ to an influx of $49.1 \text{ mmol m}^{-2} \text{ d}^{-1}$. Buffalo Pound Lake also exhibited ingassing of CO_2 in summer or fall seasons of four years (1979, 1987, 1991, and 2012), but none of these events resulted in the basin experiencing a net influx of CO_2 when calculated at an annual scale. Over the entire 36-yr dataset, total spring CO_2 efflux averaged 63.8% of total flux ($\pm 19.8\%$), but this value declines to 32.9% ($\pm 19.8\%$) when only open-water spring flux is considered. In contrast, CO_2 efflux was lowest in summer (14.0% of annual total) and increased slightly in fall (22.2%).

There were no pronounced decadal-scale trends in the estimated CO_2 content or effluxes from Buffalo Pound Lake (Fig. 2). Seasonal averages of $p\text{CO}_2$ (μatm) varied by year, with spring $p\text{CO}_2$ being the highest, averaging $1797 \mu\text{atm}$ (range $480.1\text{--}3334 \mu\text{atm}$), summer $p\text{CO}_2$ is the lowest and with less variability (average 683 , range $62.3\text{--}1387 \mu\text{atm}$), and fall intermediate between the spring and summer (average 818.9 , range $327\text{--}1981 \mu\text{atm}$). Winter $p\text{CO}_2$ averaged $1730 \mu\text{atm}$ with a minimum of 499 and a maximum of 3687 . We did not see

any significant ($p > 0.1$) decadal-scale trends with $p\text{CO}_2$ in each season vs. year (regression of data in Fig. 2) or with annual $p\text{CO}_2$ averages vs. year.

The best-fitting GAM to model the $p\text{CO}_2$ time series was a tensor-product smooth of sampling date and day of year as a seasonal trend, which varied smoothly with the between-year trend. This model was also the most complex in terms of the effective degrees of freedom ($\text{EDF} = 506.8$), but provided better fit to observed data ($\text{AIC} = 25,769$) than did the next best model ($\text{AIC} = 25,926$). The best-fit model explained 97% of deviance in the $p\text{CO}_2$ data, with an adjusted R^2 of 0.90. Qualitatively, the best-fit model also better explained large annual peaks in $p\text{CO}_2$, as well as year-to-year variation in the magnitude of that peak. Overall, the mean $p\text{CO}_2$ estimated by the best-fit model oscillated slowly over the 36 yr of study and did not exhibit sudden changes between years.

In most years, modeled $p\text{CO}_2$ increased under ice, declined substantially at spring ice melt, stayed low during summer, and, with a few exceptions, remained low until ice formation in the fall (Fig. 3). Ice-cover duration varied > 7 weeks across the 36-yr period, from a minimum of 133 d in 2000 to a maximum of 183 d in 1979 (mean 156.7 ± 12.5 d). The length of spring CO_2 decline varied from 3 weeks (in 1986 and 1988) to 15 weeks (1991) with a mean ($\pm \text{SD}$) of $10.5 (\pm 2.9)$ weeks. On average, the spring $p\text{CO}_2$ decline started 4.9 ± 2.3 weeks before the observed date of ice melt. In contrast, there were few indications of a sudden change in estimated $p\text{CO}_2$ in fall such as would be expected if CO_2 was released suddenly from hypolimnetic waters at fall mixis (Vachon and del Giorgio 2014; Ducharme-Riel et al. 2015). Instead, $p\text{CO}_2$ peaked during winter in most years (Fig. 3b), with only a few years showing limited CO_2 build-up under ice (1984, 1987, 1995, and 2012). There were no statistically significant trends in the relationship between fall $p\text{CO}_2$ and the timing of ice formation.

Comparison of GAMs developed independently for spring, summer, and fall seasons revealed that the influence of antecedent seasonal conditions on $p\text{CO}_2$ declined from spring to fall (Fig. 4; Supporting Information). For example, the best-fit GAM for spring $p\text{CO}_2$ used the scaled t distribution for heavily tailed data and explained 72.6% of the deviance ($R^2_{\text{adj}} = 0.64$, $n = 31$). This model demonstrated that mean values increased with the previous summer's average Chl a concentrations and the duration of ice cover but declined with spring water temperature (Fig. 4A–C) and was comparable when only open-water spring $p\text{CO}_2$ was considered (77.5% deviance explained, data not shown). In contrast, GAMs were significant but less predictive for both mean summer and fall $p\text{CO}_2$ values. In these models, mean summer $p\text{CO}_2$ declined with increases in mean summer Chl a concentrations but tended to increase with $p\text{CO}_2$ recorded in the previous spring, particularly at high values (Fig. 4D,E; 43.6% deviance explained, $R^2_{\text{adj}} = 0.4$, $n = 36$). Similarly, fall




Fig. 3. (a) GAM of $p\text{CO}_2$ by year and day of year in Buffalo Pound based on 36 yr of weekly data. Line color represents year of sampling and shows little evidence of directional change in lake $p\text{CO}_2$ in the 36-yr sampling period. **(b)** GAM model as represented in **(a)** but separated by year. Vertical lines indicate ice-off and ice-on dates. Most years demonstrate a consistent pattern of increasing $p\text{CO}_2$ under ice in winter, followed by efflux in spring, low $p\text{CO}_2$ in summer and fall, and then CO_2 accumulation under ice in fall.

$p\text{CO}_2$ declined with elevated autumnal Chl *a* concentrations but showed more complex (unimodal) relationships with both summer $p\text{CO}_2$ and water temperature during fall

(Fig. 4F–H; 23.3% deviance explained, $R^2_{\text{adj}} = 0.28$, $n = 34$). DOC was not a significant predictor and did not end up in any final model.




Fig. 4. GAM results for seasonally averaged $p\text{CO}_2$ in Buffalo Pound. Spring average $p\text{CO}_2$ was best explained with a combination of **(a)** Chl a concentration in the previous summer, **(b)** ice-cover duration, and **(c)** current water temperature (GAM deviance explained = 72.6%), while summer average $p\text{CO}_2$ was explained using **(d)** current summer Chl a concentration and **(e)** average $p\text{CO}_2$ in the preceding spring (deviance explained = 43.6%), and fall was best explained using **(f)** current fall Chl a concentration, **(g)** preceding summer $p\text{CO}_2$, and **(h)** current fall water temperature (deviance explained = 23.3%). Plots are partial plots of the smooth terms in the model, and the y axis is the intercept plus the partial effect of the individual smooths.

Discussion

Weekly estimates of CO_2 content of Buffalo Pound Lake over 36 yr demonstrated that a eutrophic hard-water lake could

remain a net source of CO_2 to the atmosphere (Fig. 1) despite elevated algal production and pH (Finlay et al. 2009, 2015). However, we found clear evidence of legacy effects in all seasons

(Fig. 4). Specifically, GAMs suggested that spring CO_2 content and efflux rates were derived mainly from winter metabolism of organic matter that was produced the previous summer, with longer ice-cover duration serving to increase both factors (Kratz et al. 1987; Baehr and DeGrandpre 2004). While CO_2 effluxes were lower during summer and fall than in spring (Fig. 3), $p\text{CO}_2$ values in summer and fall were also regulated by interactions between current lake production and the legacy of the previous season's CO_2 concentration. Overall, we found no evidence of a major release of CO_2 during fall, as often occurs in thermally stratified lakes (Hesslein et al. 1991; Cole et al. 1994; Ducharme-Riel et al. 2015) and conclude that annual CO_2 budgets are strongly influenced by spring efflux, and therefore antecedent limnological conditions (Fig. 4). Given the highly synchronous patterns in CO_2 and other limnological parameters of lakes in this region (Vogt et al. 2011; Finlay et al. 2015), we believe that these results will be representative of other lakes in the Northern Great Plains (e.g., Meding and Jackson 2003; Donald et al. 2015; Maheaux et al. 2016).

Temporal variation in CO_2

Total CO_2 flux in spring strongly influenced the magnitude of annual CO_2 flux in Buffalo Pound, owing to the elevated respiratory-derived CO_2 under ice. Total spring efflux of CO_2 averaged $9.69 \text{ mol } \text{CO}_2 \text{ m}^{-2}$, which is comparable to that observed in DOC-rich Wisconsin and Finnish lakes ($1.1\text{--}13.7 \text{ mmol } \text{CO}_2 \text{ m}^{-2} \text{ spring}^{-1}$; Striegl et al. 2001), whereas vernal open-water release alone was $32.9\% \pm 19.8\%$ of annual values. Summer $p\text{CO}_2$ remained relatively low and, with the exception of a few anomalous years, there was little indication of hypolimnetic CO_2 release in fall (Vachon and del Giorgio 2014; Ducharme-Riel et al. 2015). Mean fall values for $p\text{CO}_2$ and CO_2 flux were only slightly higher than in summer, consistent with the polymictic status of Buffalo Pound and the irregular occurrence of thermal stratification during summer (Dröscher et al. 2009). In general, Buffalo Pound Lake was a net annual source of CO_2 to the atmosphere, with mean total annual flux of CO_2 ($18.53 \text{ mol } \text{m}^{-2} \text{ yr}^{-1}$) and mean daily flux rates ($103.95 \text{ mmol } \text{m}^{-2} \text{ d}^{-1}$), comparable to some other hard-water systems (Striegl and Michmerhuizen 1998) but higher than boreal lakes (Rantakari and Kortelainen 2005; Abnizova et al. 2012; Ducharme-Riel et al. 2015).

Atmospheric exchange of CO_2 during spring accounted for 63.8% of annual CO_2 efflux from Buffalo Pound Lake, a value much higher than that found elsewhere (Anderson et al. 1999; Ducharme-Riel et al. 2015) and reported earlier for this site (Finlay et al. 2015). However, this result includes both the "potential" and "open-water" spring fluxes and assumes both that ice is highly permeable to gas exchange prior to its complete disappearance and that CO_2 efflux to the atmosphere is the main mechanism reducing CO_2 content during spring. If instead, gas exchange is limited through even fully fractured ice due to limited hydrologic and atmospheric exchange (Loose et al. 2011) and degassing occurs following formation of marginal (lateral)

open water immediately prior to full ice melt (Loose and Schlosser 2011), then CO_2 efflux declines to $\sim 33\%$ of annual flux, a value more in line with boreal systems. Furthermore, we note that CO_2 could have declined under ice in Buffalo Pound due to alternate mechanisms, including elevated primary production by attached and motile algae (Salmi and Salonen 2016; Hampton et al. 2017), redistribution of CO_2 -rich deep waters by convective water-column currents (Kelley 1997; Mironov et al. 2002; Pernica et al. 2017) or chemical dissolution of sedimentary CaCO_3 (Finlay et al. 2015). Taken together, these observations suggest that our spring estimates represent the maximum possible CO_2 efflux and illustrate that intensive studies of under-ice processes in the weeks prior to ice melt are needed to fully characterize the magnitude and importance of vernal CO_2 release.

Here, our GAM analyses reaffirmed that warmer winters with reduced ice-cover duration results in lower winter CO_2 accumulation and thus emissions in spring, as has been observed previously (Finlay et al. 2009, 2015). We did not, however, see similar consistent decadal trends for annual or seasonal CO_2 flux. In part, these differences arise because the duration of ice cover declined during the most recent 20-yr interval studied by Finlay et al. (2015), but not during the entire 36-yr period included here ($r^2 = -0.01$, $p = 0.5$). Given the elevated contribution of spring CO_2 fluxes to the annual budgets, as seen here and in other lake districts (Kratz et al. 1987; Cole et al. 1994; Striegl and Michmerhuizen 1998), climate change is likely to profoundly alter future lake CO_2 fluxes. Specifically, future prairie climates will be warmer and drier (Sauchyn and Kulshreshtha 2008; Lapp et al. 2009; Newton et al. 2014), with less ice cover (Shuter et al. 2013), patterns that should reduce the magnitude of CO_2 emissions in spring. This reduced spring CO_2 flux will translate into a reduction of long-term annual flux provided there is an alternative loss pathway for this C, such as carbonate precipitation or organic matter sedimentation (Tranvik et al. 2009), both of which are common in productive, hard-water lakes. Although reduced ice cover can potentially affect annual primary production, we found no significant relationship between ice duration and summer Chl *a* ($r^2 = 0.061$, $p = 0.15$) during the past 36 yr when ice cover varied from 133 to 183 d. Instead, given that regional phytoplankton biomass is a complex function of temperature and nutrient influx (Vogt et al. 2018) and given the legacy effects seen herein, we infer that further changes in spring and annual CO_2 emissions will also depend heavily on the effectiveness of nutrient management strategies (Leavitt et al. 2006; Bunting et al. 2016).

Controls of seasonal $p\text{CO}_2$

Controls of CO_2 content and potential atmospheric exchange were strongly influenced by lake production but differed in form and function among seasons. During spring, $p\text{CO}_2$ was strongly and positively influenced by mean Chl *a* content of the previous summer, consistent with microbial respiration of autochthonous

organic matter consuming oxygen (Meding and Jackson 2003; Powers et al. 2017) and producing CO_2 (Kratz et al. 1987; Finlay et al. 2015) under ice. While $p\text{CO}_2$ levels in spring were also enhanced by the duration of ice cover (longer time for CO_2 accumulation) and cool spring temperatures (enhanced gas solubility), the paramount effect of Chl *a* may reflect the highly eutrophic conditions in Buffalo Pound (Hall et al. 1999; McGowan et al. 2005). Although present at elevated concentrations, allochthonous DOC in Buffalo Pound is recalcitrant relative to other sources (Williamson et al. 1999; Guillemette et al. 2017), largely unrelated to rates of microbial production (Finlay et al. 2010), and was not included in our final models of CO_2 content. Instead, it appears that factors regulating mid-summer production may be unanticipated but important controls of subsequent spring CO_2 efflux.

GAMs also suggested that coeval Chl *a* was correlated negatively with $p\text{CO}_2$ during in both summer and fall, consistent with a strong role of phytoplankton uptake during photosynthesis as seen in other autotrophic lakes (del Giorgio and Peters 1994). Additional influences of microbial processes are indicated also by the presence of a positive relationship between temperature and $p\text{CO}_2$ in fall, a pattern consistent with the role of bacterial respiration of OM (del Giorgio and Peters 1994; Cole et al. 1994), rather than changes in gas solubility as the lake cools (Pinho et al. 2016). Again, we were unable to detect an effect of DOC on $p\text{CO}_2$ from Buffalo Pound in either summer or fall models, possibly because groundwater inputs, carbonate buffering, calcification, and anaerobic metabolism also decouple the relationship between allochthonous DOC influx and microbial metabolism in regional lakes (Bogard and del Giorgio 2016; Stets et al. 2017). Full carbon budgets in each season, including catchment loading of organic and inorganic carbon, would be required in order to fully evaluate these alternatives.

Legacy effects of climate and limnological conditions

Analysis of a 36-yr continuous time series of water chemistry demonstrated that instantaneous estimates of CO_2 content in lake waters are regulated by current limnological conditions as well as the persistent influence of lake conditions in earlier seasons. Such legacy effects are well known from studies of terrestrial biogeochemistry (Cuddington 2011) and land-water linkages (Martin et al. 2011) but are less well understood for in situ biogeochemical cycles (Meding and Jackson 2003; Hampton 2015). Recent studies suggest that spring water chemistry is strongly influenced by under-ice processes (Powers et al. 2017), and, consistent with that view, our GAM analysis showed that vernal $p\text{CO}_2$ could be predicted best (72.6% of deviance explained) from a combination of mean Chl *a* concentration during the previous summer, ice-cover duration during the antecedent winter, and coeval spring water temperature. Overall, historical parameters had a paramount effect on spring model performance (Fig. 4), while coeval biological parameters were nonsignificant (e.g., spring Chl *a*),

suggesting that vernal CO_2 fluxes were controlled mainly by limnological and climatic conditions in earlier seasons. Given the potential importance of spring CO_2 efflux to the annual CO_2 budgets (see above), these findings suggest that atmospheric warming (Finlay et al. 2015) and surface water eutrophication (Leavitt et al. 2006; Bunting et al. 2016) will interact in complex manners to regulate the importance of hard-water lakes in global carbon budgets.

Comparison of GAMs developed for individual seasons demonstrates that the strength of legacy effects declines continuously through the ice-free period, both in terms of explanatory power and the influence of historical parameters on gas fluxes (Fig. 4). Specifically, the predictive power of GAMs declined by ~ 50% in successive seasons, from spring (72.6%) to summer (43.6%) and fall (23.3%), while only $p\text{CO}_2$ levels during the preceding season were retained as a secondary predictor in summer and fall models. Although speculative, we infer that these declines may reflect the progressive accumulation of effects of intervening meteorological events (e.g., wind and low pressure cells), which are known to influence CO_2 fluxes at the scale of days to weeks (Morales-Pineda et al. 2014). In addition, as coeval Chl *a* concentrations were retained in latter models, yet are strongly influenced by summer water temperatures and nutrient content in Buffalo Pound and other regional lakes (Vogt et al. 2018), we infer that immediate controls of these limnological parameters may override the importance of legacy effects on CO_2 content and flux.

Conclusion

These analyses used a 36-yr time series with weekly resolution from a hard-water lake to demonstrate that instantaneous $p\text{CO}_2$ is regulated by a combination of current limnological conditions and legacy effects from earlier seasons. This legacy effect was most pronounced in spring and declined throughout the ice-free season. The form and identity of physicochemical controls also changed through time, with climate (ice-cover duration) being the strongest predictor in spring model, and coeval estimates of lake metabolism (Chl *a*) mainly regulating $p\text{CO}_2$ in summer and fall. The strength of these relationships reflects our ability to predict CO_2 in the future. Specifically, the strong spring relationship suggests that future climate warming and reduction of ice cover will diminish the importance of antecedent conditions, and may reduce annual CO_2 emissions to the atmosphere, particularly if efforts to reduce nutrient loading in this region are successful. This relationship explained 72% of the variability in the data, which allows for predictions of how future climate change will affect lake carbon processing in this system.

References

- Abnizova, A., J. Siemens, M. Langer, and J. Boike. 2012. Small ponds with major impact: The relevance of ponds

- and lakes in permafrost landscapes to carbon dioxide emissions. *Global Biogeochem. Cycles* **26**: GB2041. doi:[10.1029/2011GB004237](https://doi.org/10.1029/2011GB004237)
- Anderson, D. E., R. G. Striegl, D. I. Stannard, C. M. Michmerhuizen, T. A. McConnaughey, and J. W. Labaugh. 1999. Estimating lake-atmosphere CO₂ exchange. *Limnol. Oceanogr.* **44**: 988–1001. doi:[10.4319/lo.1999.44.4.0988](https://doi.org/10.4319/lo.1999.44.4.0988)
- Baehr, M. M., and M. D. Degrandpre. 2002. Under-ice CO₂ and O₂ variability in a freshwater lake. *Biogeochemistry* **61**: 95–113. doi:[10.1023/A:1020265315833](https://doi.org/10.1023/A:1020265315833)
- Baehr, M. M., and M. D. Degrandpre. 2004. In situ pCO₂ and O₂ measurements in a lake during turnover and stratification: Observations and modeling. *Limnol. Oceanogr.* **49**: 330–340. doi:[10.4319/lo.2004.49.2.0330](https://doi.org/10.4319/lo.2004.49.2.0330)
- Bogard, M. J., and P. A. del Giorgio. 2016. The role of metabolism in modulating CO₂ fluxes in boreal lakes. *Global Biogeochem. Cycles* **30**: 1509–1525. doi:[10.1002/2016GB005463](https://doi.org/10.1002/2016GB005463)
- Bunting, L., P. R. Leavitt, G. L. Simpson, B. Wissel, K. R. Laird, B. F. Cumming, A. S. Amand, and D. R. Engstrom. 2016. Increased variability and sudden ecosystem state change in Lake Winnipeg, Canada, caused by 20th century agriculture. *Limnol. Oceanogr.* **61**: 2090–2107. doi:[10.1002/lno.10355](https://doi.org/10.1002/lno.10355)
- Cole, J. J., N. F. Caraco, G. W. Kling, and T. K. Kratz. 1994. Carbon dioxide supersaturation in the surface waters of lakes. *Science* **265**: 1569–1570. doi:[10.1126/science.265.5178.1568](https://doi.org/10.1126/science.265.5178.1568)
- Cole, J. J., and others. 2007. Plumbing the global carbon cycle: Integrating inland waters into the terrestrial carbon budget. *Ecosystems* **10**: 172–185. doi:[10.1007/s10021-006-9013-8](https://doi.org/10.1007/s10021-006-9013-8)
- Cuddington, K. 2011. Legacy effects: The persistent impact of ecological interactions. *Biol. Theory* **6**: 203–210. doi:[10.1007/s13752-012-0027-5](https://doi.org/10.1007/s13752-012-0027-5)
- del Giorgio, P. A., and R. H. Peters. 1994. Patterns in planktonic P:R ratios in lakes: Influence of lake trophicity and dissolved organic carbon. *Limnol. Oceanogr.* **39**: 772–787. doi:[10.4319/lo.1994.39.4.0772](https://doi.org/10.4319/lo.1994.39.4.0772)
- Denfeld, B. A., P. Kortelainen, M. Rantakari, S. Sobek, and G. A. Weyhenmeyer. 2015. Regional variability and drivers of below ice CO₂ in boreal and subarctic lakes. *Ecosystems* **19**: 461–476. doi:[10.1007/s10021-015-9944-z](https://doi.org/10.1007/s10021-015-9944-z)
- Dillon, P. J., and L. A. Molot. 1997. Dissolved organic and inorganic carbon mass balances in central Ontario lakes. *Biogeochemistry* **36**: 29–42. doi:[10.1023/A:1005731828660](https://doi.org/10.1023/A:1005731828660)
- Donald, D. B., B. R. Parker, J.-M. Davies, and P. R. Leavitt. 2015. Nutrient sequestration in the Lake Winnipeg watershed. *J. Great Lakes Res.* **41**: 630–642. doi:[10.1016/j.jglr.2015.03.007](https://doi.org/10.1016/j.jglr.2015.03.007)
- Dröschner, I., A. Patoine, K. Finlay, and P. R. Leavitt. 2009. Climate control of the spring clear-water phase through the transfer of energy and mass to lakes. *Limnol. Oceanogr.* **54**: 2469–2480. doi:[10.4319/lo.2009.54.6_part_2.2469](https://doi.org/10.4319/lo.2009.54.6_part_2.2469)
- Ducharme-Riel, V., D. Vachon, P. A. del Giorgio, and Y. T. Prairie. 2015. The relative contribution of winter under-ice and summer hypolimnetic CO₂ accumulation to the annual CO₂ emissions from northern lakes. *Ecosystems* **18**: 1–13. doi:[10.1007/s10021-015-9846-0](https://doi.org/10.1007/s10021-015-9846-0)
- Finlay, K., P. R. Leavitt, B. Wissel, and Y. T. Prairie. 2009. Regulation of spatial and temporal variability of carbon flux in six hard-water lakes of the northern Great Plains. *Limnol. Oceanogr.* **54**: 2553–2564. doi:[10.4319/lo.2009.54.6_part_2.2553](https://doi.org/10.4319/lo.2009.54.6_part_2.2553)
- Finlay, K., P. R. Leavitt, A. Patoine, and B. Wissel. 2010. Magnitudes and controls of organic and inorganic carbon flux through a chain of hard-water lakes on the northern Great Plains. *Limnol. Oceanogr.* **55**: 1551–1564. doi:[10.4319/lo.2010.55.4.1551](https://doi.org/10.4319/lo.2010.55.4.1551)
- Finlay, K., R. J. Vogt, M. J. Bogard, B. Wissel, B. M. Tutolo, G. L. Simpson, and P. R. Leavitt. 2015. Decrease in CO₂ efflux from northern hardwater lakes with increasing atmospheric warming. *Nature* **519**: 1–13. doi:[10.1038/nature14172](https://doi.org/10.1038/nature14172)
- Gerten, D., and R. Adrian. 2000. Climate-driven changes in spring plankton dynamics and the sensitivity of shallow polymictic lakes to the North Atlantic Oscillation. *Limnol. Oceanogr.* **45**: 1058–1066. doi:[10.4319/lo.2000.45.5.1058](https://doi.org/10.4319/lo.2000.45.5.1058)
- Guillemette, F. O., E. von Wachenfeldt, D. N. Kothawala, D. Bastviken, and L. J. Tranvik. 2017. Preferential sequestration of terrestrial organic matter in boreal lake sediments. *J. Geophys. Res. Biogeosci.* **122**: 863–874. doi:[10.1002/2016JG003735](https://doi.org/10.1002/2016JG003735)
- Hall, R. I., P. R. Leavitt, and A. S. Dixit. 1999. Limnological succession in reservoirs: A paleolimnological comparison of two methods of reservoir formation. *Can. J. Fish. Aquat. Sci.* **56**: 1109–1121. doi:[10.1139/f99-047](https://doi.org/10.1139/f99-047)
- Hammer, U. T. 1986. Saline lake ecosystems of the world. Dr. W. Junk. doi:[10.1016/0011-2240\(86\)90062-3](https://doi.org/10.1016/0011-2240(86)90062-3)
- Hampton, S. E., and others. 2017. Ecology under lake ice. *Ecology Lett.* **20**: 98–111. doi:[10.1111/ele.12699](https://doi.org/10.1111/ele.12699)
- Hastie, T. J., and R. J. Tibshirani. 1990. Generalized Additive Models, Chapman & Hall / CRC. Boca Raton.
- Hesslein, R. H., J. W. M. Rudd, C. Kelly, P. Ramlal, and K. A. Hallard. 1991. Carbon dioxide pressure in surface waters of Canadian lakes, p. 413–443. *In* S. C. Wilhelms and J. S. Gulliver [eds.], Air-water mass transfer. American Society of Civil Engineers.
- Hoover, T. E., and D. C. Berkshire. 1969. Effects of hydration on carbon dioxide exchange across an air-water interface. *J. Geophys. Res.* **74**: 456–464. doi:[10.1029/JB074i002p00456](https://doi.org/10.1029/JB074i002p00456)
- Kelley, D. E. 1997. Convection in ice-covered lakes: Effects of algal suspension. *J. Plankton Res.* **19**: 1859–1880. doi:[10.1093/plankt/19.12.1859](https://doi.org/10.1093/plankt/19.12.1859)
- Kratz, T. K., R. B. Cook, and C. J. Bowser. 1987. Winter and spring pH depressions in northern Wisconsin lakes caused by increases in pCO₂. *Can. J. Fish. Aquat. Sci.* **44**: 1082–1088. doi:[10.1139/f87-129](https://doi.org/10.1139/f87-129)
- Lapp, S., D. Sauchyn, and B. Toth. 2009. Constructing scenarios of future climate and water supply for the SSRB: Use and limitations for vulnerability assessment. *Prairie Forum* **34**: 153–180.
- Leavitt, P. R., C. S. Brock, C. Ebel, and A. Patoine. 2006. Landscape-scale effects of urban nitrogen on a chain of

- freshwater lakes in central North America. *Limnol. Oceanogr.* **51**: 2262–2277. doi:[10.4319/lo.2006.51.5.2262](https://doi.org/10.4319/lo.2006.51.5.2262)
- Loose, B., and P. Schlosser. 2011. Sea ice and its effect on CO₂ flux between the atmosphere and the Southern Ocean interior. *J. Geophys. Res. Oceans* **116**: C11019. doi:[10.1029/2010JC006509](https://doi.org/10.1029/2010JC006509)
- Loose, B., and others. 2011. Gas diffusion through columnar laboratory sea ice: Implications for mixed-layer ventilation of CO₂ in the seasonal ice zone. *Tellus B Chem. Phys. Meteorol.* **63**: 23–39. doi:[10.1111/j.1600-0889.2010.00506.x](https://doi.org/10.1111/j.1600-0889.2010.00506.x)
- López Bellido, J., T. Tulonen, P. Kankaala, and A. Ojala. 2012. Concentrations of CO₂ and CH₄ in water columns of two stratified boreal lakes during a year of atypical summer precipitation. *Biogeochemistry* **113**: 613–627. doi:[10.1007/s10533-012-9792-2](https://doi.org/10.1007/s10533-012-9792-2)
- Maberly, S. C. 1996. Diel, episodic and seasonal changes in pH and concentrations of inorganic carbon in a productive lake. *Freshwat. Biol.* **35**: 579–598. doi:[10.1111/j.1365-2427.1996.tb01770.x](https://doi.org/10.1111/j.1365-2427.1996.tb01770.x)
- Maheaux, H., P. R. Leavitt, and L. J. Jackson. 2016. Asynchronous onset of eutrophication among shallow prairie lakes of the northern Great Plains, Alberta, Canada. *Glob. Chang. Biol.* **22**: 271–283. doi:[10.1111/gcb.13076](https://doi.org/10.1111/gcb.13076)
- Martin, S. L., D. B. Hayes, D. T. Rutledge, and D. W. Hyndman. 2011. The land-use legacy effect: Adding temporal context to lake chemistry. *Limnol. Oceanogr.* **56**: 2362–2370. doi:[10.4319/lo.2011.56.6.2362](https://doi.org/10.4319/lo.2011.56.6.2362)
- McGowan, S., P. R. Leavitt, and R. I. Hall. 2005. A whole-lake experiment to determine the effects of winter droughts on shallow lakes. *Ecosystems* **7**: 694–708. doi:[10.1007/s10021-005-0152-x](https://doi.org/10.1007/s10021-005-0152-x)
- Meding, M. E., and L. J. Jackson. 2003. Biotic, chemical and morphometric factors contributing to winter anoxia in prairie lakes. *Limnol. Oceanogr.* **48**: 1633–1642. doi:[10.4319/lo.2003.48.4.1633](https://doi.org/10.4319/lo.2003.48.4.1633)
- Mironov, D., A. Terzhevik, G. Kirillin, T. Jonas, J. Malm, and D. Farmer. 2002. Radiatively driven convection in ice-covered lakes: Observations, scaling, and a mixed layer model. *J. Geophys. Res.* **107**: 3032. doi:[10.1029/2001JC000892](https://doi.org/10.1029/2001JC000892)
- Morales-Pineda, M., A. Cózar, I. Laiz, B. Úbeda, and J. Á. Gálvez. 2014. Daily, biweekly, and seasonal temporal scales of pCO₂ variability in two stratified Mediterranean reservoirs. *J. Geophys. Res. Biogeosci.* **119**: 509–520. doi:<https://doi.org/10.1007/s10021-003-0152-x>
- Newton, B. W., T. D. Prowse, and B. R. Bonsal. 2014. Evaluating the distribution of water resources in western Canada using synoptic climatology and selected teleconnections. Part 2: Summer season. *Hydrol. Processes* **28**: 4235–4249. doi:[10.1002/hyp.10235](https://doi.org/10.1002/hyp.10235)
- Pernica, P., R. L. North, and H. M. Baulch. 2017. In the cold light of day: The potential importance of under-ice convective mixed layers to primary producers. *Inland Wat.* **7**: 138–150. doi:[10.1080/20442041.2017.1296627](https://doi.org/10.1080/20442041.2017.1296627)
- Pinho, L., C. M. Duarte, H. Marotta, and A. Enrich-Prast. 2016. Temperature dependence of the relationship between pCO₂ and dissolved organic carbon in lakes. *Biogeosciences* **13**: 865–871. doi:[10.5194/bg-13-865-2016](https://doi.org/10.5194/bg-13-865-2016)
- Powers, S. M., S. G. Labou, H. M. Baulch, R. J. Hunt, N. R. Lottig, S. E. Hampton, and E. H. Stanley. 2017. Ice duration drives winter nitrate accumulation in north temperate lakes. *Limnol. Oceanogr. Lett.* **2**: 177–189. doi:[10.1002/lo2.10048](https://doi.org/10.1002/lo2.10048)
- Prairie, Y. T. 2008. Carbocentric limnology: Looking back, looking forward. *Can. J. Fish. Aquat. Sci.* **65**: 543–548. doi:[10.1139/f08-011](https://doi.org/10.1139/f08-011)
- R Core Team. 2018. R: A language and environment for statistical computing. R Foundation for Statistical Computing. Available from <https://www.R-project.org/>
- Rantakari, M., and P. Kortelainen. 2005. Interannual variation and climatic regulation of the CO₂ emission from large boreal lakes. *Glob. Chang. Biol.* **11**: 1368–1380. doi:[10.1111/j.1365-2486.2005.00982.x](https://doi.org/10.1111/j.1365-2486.2005.00982.x)
- Salmi, P., and K. Salonen. 2016. Regular build-up of the spring phytoplankton maximum before ice-break in a boreal lake. *Limnol. Oceanogr.* **61**: 240–253. doi:[10.1002/lno.10214](https://doi.org/10.1002/lno.10214)
- Sauchyn, D., and S. Kulshreshtha. 2008. The Prairies. From impacts to adaptation: Canada in a changing climate. Government of Canada. Cat. No.: M174-2/1-2007
- Shuter, B. J., C. K. Minns, and S. R. Fung. 2013. Empirical models for forecasting changes in the phenology of ice cover for Canadian lakes. *Can. J. Fish. Aquat. Sci.* **70**: 982–991. doi:[10.1139/cjfas-2012-0437](https://doi.org/10.1139/cjfas-2012-0437)
- Simpson, G. L. 2018. Modelling Palaeoecological Time Series Using Generalised Additive Models. *Frontiers in Ecology and Evolution* **6**: 149. doi:[10.3389/fevo.2018.00149](https://doi.org/10.3389/fevo.2018.00149)
- Stets, E. G., R. G. Striegl, G. R. Aiken, D. O. Rosenberry, and T. C. Winter. 2009. Hydrologic support of carbon dioxide flux revealed by whole-lake carbon budgets. *J. Geophys. Res.* **114**: G01008. doi:[10.1029/2008JG000783](https://doi.org/10.1029/2008JG000783)
- Stets, E. G., D. Butman, C. P. McDonald, S. M. Stackpoole, M. D. Degrandpre, and R. G. Striegl. 2017. Carbonate buffering and metabolic controls on carbon dioxide in rivers. *Global Biogeochem. Cycles* **31**: 663–677. doi:[10.1002/2016GB005578](https://doi.org/10.1002/2016GB005578)
- Striegl, R. G., and C. M. Michmerhuizen. 1998. Hydrologic influence on methane and carbon dioxide dynamics at two north-Central Minnesota lakes. *Limnol. Oceanogr.* **43**: 1519–1529. doi:[10.4319/lo.1998.43.7.1519](https://doi.org/10.4319/lo.1998.43.7.1519)
- Striegl, R. G., P. Kortelainen, J. P. Chanton, K. P. Wickland, G. C. Bugna, and M. Rantakari. 2001. Carbon dioxide partial pressure and ¹³C content of north temperate and boreal lakes at spring ice melt. *Limnol. Oceanogr.* **46**: 941–945. doi:[10.4319/lo.2001.46.4.0941](https://doi.org/10.4319/lo.2001.46.4.0941)
- Tranvik, L. J., and others. 2009. Lakes and reservoirs as regulators of carbon cycling and climate. *Limnol. Oceanogr.* **54**: 2298–2314. doi:[10.4319/lo.2009.54.6_part_2.2298](https://doi.org/10.4319/lo.2009.54.6_part_2.2298)
- Vachon, D., and Y. T. Prairie. 2013. The ecosystem size and shape dependence of gas transfer velocity versus wind

- speed relationships in lake. *Can. J. Fish. Aquat. Sci.* **70**: 1757–1764. doi:[10.1139/cjfas-2013-0241](https://doi.org/10.1139/cjfas-2013-0241)
- Vachon, D., and P. A. del Giorgio. 2014. Whole-Lake CO_2 dynamics in response to storm events in two morphologically different lakes. *Ecosystems* **17**: 1338–1353. doi:[10.1007/s10021-014-9799-8](https://doi.org/10.1007/s10021-014-9799-8)
- Vogt, R. J., J. A. Rusak, A. Patoine, and P. R. Leavitt. 2011. Differential effects of energy and mass influx on the landscape synchrony of lake ecosystems. *Ecology* **92**: 1104–1114. doi:[10.1890/10-1846.1](https://doi.org/10.1890/10-1846.1)
- Vogt, R. J., S. Sharma, and P. R. Leavitt. 2018. Direct and interactive effects of climate, meteorology, river hydrology, and lake characteristics on water quality in productive lakes of the Canadian Prairies. *Can. J. Fish. Aquat. Sci.* **75**: 47–59. doi:[10.1139/cjfas-2016-0520](https://doi.org/10.1139/cjfas-2016-0520)
- Wetzel, R. G. 2001. *Limnology*. Academic Press.
- Wickham, H. 2009. *ggplot2: Elegant graphics for data analysis*. Springer-Verlag.
- Williamson, C. E., D. P. Morris, M. L. Pace, and O. G. Olson. 1999. Dissolved organic carbon and nutrients as regulators of lake ecosystems: Resurrection of a more integrated paradigm. *Limnol. Oceanogr.* **44**: 795–803. doi:[10.4319/lo.1999.44.3_part_2.0795](https://doi.org/10.4319/lo.1999.44.3_part_2.0795)
- Winder, M., and D. E. Schindler. 2004. Climatic effects on the phenology of lake processes. *Glob. Chang. Biol.* **10**: 1844–1856. doi:[10.1111/j.1365-2486.2004.00849.x](https://doi.org/10.1111/j.1365-2486.2004.00849.x)
- Wood, S. N. 2017. *Generalized additive models: An introduction with R*, 2nd ed. CRC Press.
- Wood, S. N., N. Pya, and B. Säfken. 2016. Smoothing parameter and model selection for general smooth models. *J. Am. Stat. Assoc.* **111**: 1548–1563. doi:[10.1080/01621459.2016.1180986](https://doi.org/10.1080/01621459.2016.1180986)

Acknowledgments

We thank Dan Conrad from the Buffalo Pound Water Treatment Plant in Moose Jaw, Saskatchewan, for providing the water chemistry data, and Helen Baulch, University of Saskatchewan for providing the diel CO_2 data for Buffalo Pound. Emma Wiik helped to collate the data and contributed to discussions regarding the analyses. We thank K. Wickland and two anonymous reviewers for helpful comments. Funding for this project was provided by an Natural Sciences and Engineering Research Council of Canada Discovery Grants to K.F., G.L.S., and P.R.L. was also supported by the Canada Research Chair program. We acknowledge that this research was conducted on Treaty 4 territory. This is a contribution to the Qu'Appelle long-term ecological research program.

Conflict of Interest

None declared.

Submitted 30 January 2018

Revised 09 July 2018

Accepted 05 December 2018

Associate editor: Kimberly Wickland